

Table 45. Antimicrobial Resistance among *Escherichia coli* by Meat Type, 2002

	Chicken Breast (n=282)		Ground Turkey (n=304)		Ground Beef (n=295)		Pork Chop (n=184)	
Antimicrobial	# Resistant	% Resistance	# Resistant	% Resistance	# Resistant	% Resistance	# Resistant	% Resistance
TET	130	46.1%	234	77.0%	91	30.9%	97	52.7%
STR	139	49.3%	175	57.6%	28	9.5%	41	22.3%
SMX	91	32.3%	146	48.0%	29	9.8%	23	12.5%
AMP	61	21.6%	95	31.3%	18	6.1%	25	13.6%
GEN	65	23.1%	82	27.0%	1	0.3%	2	1.1%
CEP	60	21.3%	45	14.8%	17	5.8%	19	10.3%
KAN	17	6.0%	40	13.2%	7	2.4%	10	5.4%
AMC	34	12.1%	17	5.6%	6	2.0%	10	5.4%
FOX	31	11.0%	10	3.3%	4	1.4%	6	3.3%
AMI	11	3.9%	7	2.3%	4	1.4%	5	2.7%
COT	10	3.6%	12	4.0%	2	0.7%	2	1.1%
TIO	20	7.1%	3	1.0%	0	0.0%	1	0.5%
NAL	8	2.8%	13	4.3%	0	0.0%	1	0.5%
CHL	2	0.7%	1	0.3%	3	1.0%	3	1.6%
AXO	0	0.0%	0	0.0%	0	0.0%	0	0.0%
CIP	0	0.0%	0	0.0%	0	0.0%	0	0.0%